

The UNM METALS
Superfund Research
Program (SRP) conducts
field and laboratory
research to better
understand how metals in
uranium mine wastes
affect the air, land and
water and the health of
people who live nearby.

The METALS SRP is focused on mining impacts in three Native American communities in the Southwest:

- Pueblo of Laguna/ Jackpile Mine
- Red Water Pond Road Community/Northeast Church Rock Mine
- Blue Gap-Tachee
 Chapter, Navajo
 Nation/Claim 28 Mine

For information about research being conducted on the Pueblo of Laguna, contact METALS community liaison, Kyle Swimmer, at 505-262-1862, or

laguna.swimmer1680@g mail.com, or visit hsc.unm.edu/college-ofpharmacy/research-andscholarship/metals/index. html.



UNM METALS Superfund Research Center Research Brief No. 3.1


Jackpile Mine wastes source of uranium in Rio Paguate and likely source of metals building up in sediments behind Paguate Reservoir

How uranium moves and accumulates in the environment can affect the quality of surface water while also contributing to the buildup of metals in river and reservoir sediments.

Between 2014 and 2017, UNM researchers, led by geochemist Johanna Blake (now with the U.S. Geological Survey) and Professor Jose Cerrato, an environmental engineer, investigated the mobility of uranium in water and accumulation in sediments along the Rio Paguate and in the wetlands of Paguate Reservoir, located 5 kilometers (about 3.1 miles) south of the Jackpile Mine.

In a paper published in 2017 in the scientific journal, *Environmental Science: Process and Impacts*, Dr. Blake and colleagues reported results of water and sediment sampling in the Rio Paguate and Paguate Reservoir behind Mesita Dam (map below). The investigators found that ore and mine wastes on the surface of the Jackpile Mine are the source of uranium in the surface water.


They also found that the concentration, or amount, of uranium in the Rio Paguate varies seasonally, with higher concentrations during the summer monsoon season runoff. While the uranium concentration in surface water decreases downstream of the mine, the uranium itself was found to accumulate, or build up, in organic-rich sediments in the Paguate Reservoir wetland.



The UNM findings are consistent with studies conducted in the late 1970s when the Jackpile Mine was operating. New Mexico Bureau of Mines and Mineral Resources researchers found increased concentrations of uranium, other metals and radioactive elements derived from the decay of uranium in the bottom sediments of the Reservoir (Popp, et al., 1983).

These studies, spaced nearly 40 years apart, document continuing impacts of the mine on Laguna water sources – long after mining stopped and initial reclamation was conducted. They help inform discussions about the extent of remediation of the possible effects of the mine releases on downstream resources.

Sept. 2019





Historical Comparisons of Uranium Levels and Stream Flow in Rio Paguate:

The figures above, from the 2017 Blake paper (left), show how uranium concentrations increase as stream flows in the Rio Paguate increase in the summer monsoon season across two separate periods: historically, 1987 and 1993 (upper left), and recently, 2014-2016 (upper right). Virtually all uranium concentrations during runoff events in both time periods exceed the USEPA maximum contaminant level (MCL) of 30 micrograms per liter (µg/l). Maximum uranium concentrations ranged from 500 µg/l to nearly 800 µg/l in surface water at the Rio Paguate crossing (called the "ford"), identified as sample station 1 in the graphic to the right and map on the front. A uranium concentration exceeding 500 µg/l at the ford was the basis for the listing of the Jackpile Mine as a federal Superfund Site in 2013.

Citations:

Blake JM, DeVore CH, Avasarala S, Ali AM, Roldan C, Bowers F, Spilde MN, Artyushkova K, Kirk MF, Peterson E, Rodriguez-Freire L, Cerrato JM. Uranium mobility and accumulation along the Rio Paguate, Jackpile Mine in Laguna Pueblo, NM. *Environmental Science: Process and Impacts*, 2017. DOI:10.1039/c6em00612d.

Popp CJ, Hawley JW, Love DW. Radionuclide and Heavy Metal Distribution in Recent Sediments of Major Streams in the Grants Mineral Belt, N.M. Socorro: New Mexico Bureau of Mines and Mineral Resources, prepared for the U.S. Office of Surface Mining, Glll5352 (1981-82), 1983. Available at https://www.yumpu.com/en/document/view/15484953/radionuclide-and-heavy-metal-distribution-in-recent-sediments-of-

Funding: NIH/NIEHS P42 ES025589 (UNM METALS)

This material was developed in part under cited research awards to the University of New Mexico. It has not been formally reviewed by the funding agencies. The views expressed are solely those of the METALS SRP staff and do not necessarily reflect those of the agencies. The funders do not endorse any products or commercial services mentioned in this research brief. The UNM METALS Superfund Research Center is not affiliated with the USEPA's Superfund program.